


6.2 – Solving Systems of Equations Graphically

Linear-Quadratic

A Linear-Quadratic System of Equations is a linear equation and a quadratic equation involving the same two variables. The solution(s) to this system are the point(s) where the line intersects the parabola (if it does at all).

Draw pictures to represent the possible number of solutions that a linear-quadratic system can have:



Example 1 – Solve the following system of equations graphically:

- 1) $4x - y + 3 = 0$
- 2) $2x^2 + 8x - y + 3 = 0$

- a) Get the linear equation into $y = mx \pm b$ form and graph.
- b) Complete the square and graph the quadratic equation.
- c) Identify and write down the points of intersection (the solution).
- d) Verify the solution by checks.

a) ① $4x - y + 3 = 0$

$$y = 4x + 3$$

Annotations: "up" above the 4, "right" below the 4, "y-int" below the 3.

b) ② $2x^2 + 8x - y + 3 = 0$

$$y = 2x^2 + 8x + 3$$

$$y = 2(x^2 + 4x) + 3$$

$b = 4, 2, 4$


$$y = 2(x^2 + 4x + 4 - 4) + 3$$

$$y = 2(x + 2)^2 - 5$$

vertex $(-2, -5)$

$a = 2$ (double up counts)

c) $(-2, -5)$ & $(0, 3)$



<p>d) check $(-2, -5)$</p> <table border="1"> <tr> <th>LS</th> <th>RS</th> </tr> <tr> <td>$4(-2) - (-5) + 3$</td> <td>0</td> </tr> <tr> <td>$-8 + 5 + 3$</td> <td></td> </tr> <tr> <td>0</td> <td>✓</td> </tr> </table>	LS	RS	$4(-2) - (-5) + 3$	0	$-8 + 5 + 3$		0	✓	<p>check $(0, 3)$</p> <table border="1"> <tr> <th>LS</th> <th>RS</th> </tr> <tr> <td>$4(0) - (3) + 3$</td> <td>0</td> </tr> <tr> <td>$0 - 3 + 3$</td> <td></td> </tr> <tr> <td>0</td> <td>✓</td> </tr> </table>	LS	RS	$4(0) - (3) + 3$	0	$0 - 3 + 3$		0	✓
LS	RS																
$4(-2) - (-5) + 3$	0																
$-8 + 5 + 3$																	
0	✓																
LS	RS																
$4(0) - (3) + 3$	0																
$0 - 3 + 3$																	
0	✓																
<table border="1"> <tr> <th>LS</th> <th>RS</th> </tr> <tr> <td>$2(-2)^2 + 8(-2) - (-5) + 3$</td> <td>0</td> </tr> <tr> <td>$8 - 16 + 5 + 3$</td> <td></td> </tr> <tr> <td>0</td> <td>✓</td> </tr> </table>	LS	RS	$2(-2)^2 + 8(-2) - (-5) + 3$	0	$8 - 16 + 5 + 3$		0	✓	<table border="1"> <tr> <th>LS</th> <th>RS</th> </tr> <tr> <td>$2(0)^2 + 8(0) - (3) + 3$</td> <td>0</td> </tr> <tr> <td>$0 + 0 - 3 + 3$</td> <td></td> </tr> <tr> <td>0</td> <td>✓</td> </tr> </table>	LS	RS	$2(0)^2 + 8(0) - (3) + 3$	0	$0 + 0 - 3 + 3$		0	✓
LS	RS																
$2(-2)^2 + 8(-2) - (-5) + 3$	0																
$8 - 16 + 5 + 3$																	
0	✓																
LS	RS																
$2(0)^2 + 8(0) - (3) + 3$	0																
$0 + 0 - 3 + 3$																	
0	✓																

Example 2 – Is (5, 7) a solution to the system 1) $3x^2 - 10y = 5$ and 2) $-y = x - 11$?

Do a check!

$$\textcircled{1} \begin{array}{l|l} \text{LS} & \text{RS} \\ \hline 3(5^2) - 10(7) & 5 \\ 3(25) - 70 & \\ 75 - 70 & \\ 5 & \end{array}$$


$$\textcircled{2} \begin{array}{l|l} \text{LS} & \text{RS} \\ \hline -(7) & (5) - 11 \\ -7 & -6 \\ & \times \end{array}$$

(5, 7) is NOT a solution.
It is on the parabola but is not on the line

Quadratic-Quadratic

A Quadratic-Quadratic System of Equations is two quadratic equations involving the same variables. The solution(s) to this system are the point(s) where the parabola intersects the other parabola (if it does at all).

Draw pictures to represent the possible number of solutions that a quadratic-quadratic system can have:




Quadratic-Quadratic

Example 3 – Solve 1) $2x^2 - 8x + 7 - y = 0$ and 2) $y + x^2 - 4x + 2 = 0$

$$\textcircled{1} \begin{aligned} y &= (2x^2 - 8x) + 7 \\ y &= 2(x^2 - 4x) + 7 \\ b &= -4, -2, 4 \\ y &= 2(x^2 - 4x + 4 - 4) + 7 \\ y &= 2(x-2)^2 - 8 + 7 \quad \text{vertex } (2, -1) \\ y &= 2(x-2)^2 - 1 \quad a = 2 \end{aligned}$$

$$\textcircled{2} \begin{aligned} y &= (-x^2 + 4x) - 2 \\ y &= -(x^2 - 4x) - 2 \\ b &= -4, -2, 4 \\ y &= -(x^2 - 4x + 4 - 4) - 2 \quad \text{vertex } (2, 2) \\ y &= -(x-2)^2 + 4 - 2 \quad a = -1 \\ y &= -(x-2)^2 + 2 \end{aligned}$$



Solutions: (1, 1) & (3, 1)


Example 4 - Solve the system $y - x^2 + 4 = 0$ and $-2x + 2x^2 - 8 = 0$

① $y = x^2 - 4$
vertex $(0, -4)$
 $a = 1$

② $\frac{2y}{2} = \frac{2x^2 - 8}{2}$

$y = x^2 - 4$

same parabola!



∞ solutions.

6.3/6.4 – Solving Systems of Equations Algebraically

Linear-
Quadratic

For a Linear-Quadratic System of Equations, what are all the possible # of solutions?

0, 1, 2

Solutions can be found graphically, as in Section 6.2, or algebraically, using either substitution or elimination.

substitution

Example 1 – Solve the following linear-quadratic system using **substitution**:

- 1) $3x + y = -9$
- 2) $4x^2 - x + y = -9$
- a) Solve the linear equation for y .
- b) Substitute the linear equation for y in the quadratic equation.
- c) Solve the quadratic equation by factoring (if you cannot factor, use the quadratic formula).
- d) Substitute the resulting x value(s) into the original linear equation to determine the corresponding y values.

a) ① $y = -3x - 9$

b & c) ② $4x^2 - x + y = -9$

$$4x^2 - x + (-3x - 9) = -9$$

$$4x^2 - x - 3x - 9 = -9$$

$$4x^2 - 4x = 0$$

$$4x(x - 1) = 0$$

$$x = 0, 1 \quad (0, -) \quad (1, -)$$

d) ① $3x + y = -9$

$$x=0 \mid \begin{array}{l} 3(0) + y = -9 \\ y = -9 \end{array}$$

$$x=1 \mid \begin{array}{l} 3(1) + y = -9 \\ 3 + y = -9 \end{array}$$

$$y = -12$$

Solutions:

$$(0, -9) \text{ \& } (1, -12)$$

Example 2 – Solve by substitution: 1) $5x - y = 10$ and 2) $x^2 + x - 2y = 0$

① $5x - y = 10$

$$5x = 10 + y$$

$$y = 5x - 10$$

② $x^2 + x - 2y = 0$

$$x^2 + x - 2(5x - 10) = 0$$

$$x^2 + x - 10x + 20 = 0$$

$$x^2 - 9x + 20 = 0$$

$$(x - 5)(x - 4) = 0$$

$$x = 4, 5$$

$$(4, -) \quad (5, -)$$

① $5(4) - y = 10$

$$20 - y = 10$$

$$y = 10$$

$5(5) - y = 10$

$$25 - y = 10$$

$$y = 15$$

Solutions:

$$(4, 10) \text{ \& } (5, 15)$$

elimination

Now, solve the same system using **elimination**:

- 1) $5x - y = 10$
- 2) $x^2 + x - 2y = 0$

- a) Align the terms with the same degree. Since the squared term is the variable x , eliminate the y -term.
- b) Multiply one or more of the equations if necessary to have the same coefficient for y .
- c) Add or subtract the two equations to eliminate y .
- d) Solve the resulting quadratic equation by factoring or the quadratic formula to find the x coordinates of the solution(s).
- e) Substitute the resulting x value(s) into the original linear equation to determine the corresponding y values.

$\begin{array}{r} \textcircled{2} \quad x^2 + x - 2y = 0 \\ \textcircled{1} \quad (5x - y = 10) \times 2 \\ \hline x^2 + x - 2y = 0 \\ - (10x - 2y = 20) \\ \hline x^2 - 9x = -20 \end{array}$	$\begin{array}{l} x^2 - 9x + 20 = 0 \\ (x-5)(x-4) = 0 \\ x = 4, 5 \\ \hline 5(4) - y = 10 \\ y = 10 \\ \hline 5(5) - y = 10 \\ y = 15 \end{array}$
--	--

Solutions:
(4, 10) & (5, 15)

Quadratic-
Quadratic

For a Quadratic-Quadratic Systems of Equations, what are all the possible # of solutions?

0, 1, 2, ∞

Example 3 – Solve the following system first by substitution, then by elimination.

- 1) $6x^2 - x - y = -1$
- 2) $4x^2 - 4x - y = -6$

Substitution:

$$\begin{aligned} \textcircled{1} \quad y &= (6x^2 - x + 1) \\ \textcircled{2} \quad 4x^2 - 4x - y &= -6 \\ 4x^2 - 4x - (6x^2 - x + 1) &= -6 \\ 4x^2 - 4x - 6x^2 + x - 1 &= -6 \\ -2x^2 - 3x + 5 &= 0 \\ 2x^2 + 3x - 5 &= 0 \\ 2x^2 - 2x + 5x - 5 &= 0 \\ 2x(x-1) + 5(x-1) &= 0 \\ (x-1)(2x+5) &= 0 \\ x &= 1, -\frac{5}{2} \end{aligned}$$

$$\begin{aligned} \textcircled{1} \quad 6(1)^2 - 1 - y &= -1 \\ 6 - 1 + 1 &= y \\ y &= 6 \\ \textcircled{2} \quad 6\left(-\frac{5}{2}\right)^2 - \left(-\frac{5}{2}\right) - y &= -1 \\ 6\left(\frac{25}{4}\right) + \frac{5}{2} - y &= -1 \\ \frac{75}{2} + \frac{5}{2} - y &= -1 \\ \frac{75}{2} + \frac{5}{2} + \frac{2}{2} &= y \\ y &= \frac{82}{2} = 41 \end{aligned}$$


Elimination:

$$\begin{array}{r} 6x^2 - x - y = -1 \\ - (4x^2 - 4x - y = -6) \\ \hline 2x^2 + 3x = 5 \\ 2x^2 + 3x - 5 = 0 \\ \text{etc...} \end{array}$$

Solutions:
(1, 6) & $\left(-\frac{5}{2}, 41\right)$

Example 4 – A Canadian cargo plane drops a crate of emergency supplies to aid-workers on the ground. The crate drops freely at first before a parachute opens to bring the crate gently to the ground. The crate's height, h , in metres, above the ground t seconds after leaving the aircraft is given by the following two equations: $h = -4.9t^2 + 900$ represents the height of the crate during freefall. $h = -4t + 500$ represents the height of the crate with the parachute open.

- How long after the crate leaves the aircraft does the parachute open? Express your answer to the nearest hundredth of a second.
- What height above the ground is the crate when the parachute opens? Express your answer to the nearest metre.
- Verify your solution.



$$\textcircled{1} h = -4.9t^2 + 900$$

$$\textcircled{2} h = -4t + 500$$

can solve using substitution or elimination.

$$-4t + 500 = -4.9t^2 + 900$$

$$4.9t^2 - 4t - 400 = 0$$

$$4.9t^2 - 4t - 400 = 0$$

$$a = 4.9, b = -4, c = -400$$

$$t = \frac{4 \pm \sqrt{(-4)^2 - 4(4.9)(-400)}}{2(4.9)}$$

$$t = \frac{4 \pm \sqrt{7856}}{9.8}$$

$$t = 9.452, -8.636$$

↑
reject

$$h = -4t + 500$$

$$h = -4(9.452) + 500$$

$$h = 462 \text{ m}$$

The parachute opens after 9.45s,
462m from the ground.

c) Verify by doing a check for
(9.452, 462.19)

①	LS	RS
	462.19	$-4.9(9.452)^2 + 900$
		462.23

②	LS	RS
	462.19	$-4(9.452) + 500$
		462.19

* not exact due to
rounding of
solutions.

7.2A – Linear Inequalities in Two Variables

Warmup

How do we read these inequalities (from left to right)? $5 > 2$ | $-3 < -1$
5 is greater than 2 | *-3 is less than -1*

What does each symbol mean? $>$ greater than $<$ less than \geq greater than or equal to \leq less than or equal to

How do you say this aloud? $x \geq 4$
"x is greater than or equal to 4"

What are some possible answers?
4, 5, 6, 100, etc.

What is the primary difference between an **equation** and an **inequality**?
An equation has a distinct number of solutions whereas an inequality has a range (often infinite) of solutions.

Example 1 - Solve the following inequality: $3x - 7 < -5$ $x < \frac{2}{3}$
 $\frac{3x}{3} < \frac{2}{3}$

Example 2 – What are some possible answers to $-2x < 6$?
 $\frac{-2x}{-2} < \frac{6}{-2}$
 $x > -3$

How is solving an inequality like solving an equation? How is it different?
Solve an inequality just like an equation EXCEPT when multiplying or dividing both sides by a negative, FLIP the inequality

Find some solutions to $3y - 2x \geq 6$
 (0,0)? $3(0) - 2(0) \geq 6$ | $(4,3)?$ | $(1,5)$
 $0 \geq 6$ | $3(3) - 2(4) \geq 6$ | $3(5) - 2(1) \geq 6$
 \times | $1 \geq 6$ | $13 \geq 6$ ✓
There is a better way than guess + check.

There is a more efficient way to find the range of solutions for the inequality above.

steps

1. Rearrange the inequality so it's in $mx \pm b$ form. Don't forget to flip the inequality if you multiply or divide by a negative number.
2. Decide whether to use a solid line or dotted line:
 - If the inequality is \leq or \geq , points on the line are included in the solution (due to the 'equals to' line under the sign), so we keep the line solid.
 - If the inequality is $<$ or $>$, points on the line are not included in the inequality, so we draw a dotted line.
3. Graph the line using slope and y-intercept. The line is called the **boundary**.
4. For $y > mx + b$ or $y \geq mx + b$, solutions to the inequality are all of the points **above** the line, so shade above. For $y < mx + b$ or $y \leq mx + b$, shade **below** the line. The shading represents the **solution region**: all of the points that satisfy the inequality.
5. CHECK: Pick a **test point** in the shaded region and test its coordinates in the inequality. If it satisfies the inequality, you've been successful. If it doesn't satisfy the inequality, you either shaded the incorrect region, or the boundary line has been graphed incorrectly.

Example 1 – Solve the inequality by graphing $3y - 2x \geq 6$.

Get into $mx+b$ form (isolate y)

$$3y - 2x \geq 6$$

$$3y \geq 2x + 6$$


$$y \geq \frac{2}{3}x + 2$$

solid line

$$y \geq \frac{2}{3}x + 2$$

greater than so shade above

The shaded region is the set of all solutions to the inequality



CHECK: Pick one possible solution and test: ex. (0,5)

$$3(5) - 2(0) \geq 6$$

$$15 \geq 6$$

✓

***When you read an inequality for shading purposes, it must be in $y = mx + b$ form!**

Example 2 – Solve $4x - 2y > 10$. Determine if (1, 3) is part of the solution.

$$4x - 2y > 10$$

$$\frac{-2y}{-2} > \frac{-4x + 10}{-2}$$


$$y < \frac{2}{1}x - 5$$

- dashed line

- shade below

Is (1,3) in the shaded region? No, so (1,3)

is NOT a solution.



Example 3 – Solve $x \leq 4$

$x = 4$ is a vertical line

$$x \leq 4$$



Solid line

* There is no 'above' or 'below' for a vertical line so think 'left' or 'right'

$$x \leq 4$$


less than means LEFT, so shade left

check $(0, 0)$

$$x \leq 4$$

$$0 \leq 4 ?$$

✓ yes



7.2B – Systems of Linear Inequalities in Two Variables

A system of linear inequalities is: more than one linear inequality on the same graph. The solution set must satisfy each linear inequality in the system.

Example 1 – Solve the system: ① $y \geq 2x + 5$ and ② $y < -x - 5$

$$\textcircled{1} \quad y \geq 2x + 5$$


↑ ↑
Solid line

$$\textcircled{2} \quad y < -\frac{1}{2}x - 5$$

↑ ↓
dashed.

- above line ①

- below line ②



Pick one possible solution and perform a check:

$$\begin{array}{l|l} (-8, 0) \quad \textcircled{1} & 0 \geq 2(-8) + 5 \\ & 0 \geq -11 \\ & \checkmark \\ \hline & \textcircled{2} \quad 0 < -(-8) - 5 \\ & 0 < 8 - 5 \\ & 0 < 3 \quad \checkmark \end{array}$$

Example 2 – Solve the system: $3x + 2y > -6$ and $-3 \leq y \leq 3$

$$\textcircled{1} \quad 3x + 2y > -6$$

$$2y > -3x - 6$$

- above ①


$$y > -\frac{3}{2}x - 3$$

- between ②

dashed line

$$\textcircled{2} \quad -3 \leq y \leq 3$$

y is in between $y = -3$ and $y = 3$,
both solid lines



Check: $(0, 0)$

$$\begin{array}{l|l} \textcircled{1} & 3(0) + 2(0) > -6 \\ & 0 > -6 \\ & \checkmark \\ \hline \textcircled{2} & -3 \leq 0 \leq 3 \\ & \checkmark \end{array}$$

STEPS: 1. Rearrange each inequality into $mx + b$ form.

2. Graph each line, using dashed ($>$, $<$) or solid (\geq , \leq) lines.

3. To find the solution region (shaded region), look to see whether to shade above or below the first line, then above or below the second line (read the inequality in $mx + b$ form).

4. Check your solution by picking a point in your solution and testing it in each of the two original inequalities. It must satisfy both inequalities. If it doesn't, an error was made at some point, so try to find out what it is, or redo the question.

Example 3 – Solve the system of linear inequalities

$$\textcircled{1} x \geq -3, \textcircled{2} y > -4, \textcircled{3} y > 2x - 4, \textcircled{4} x + 6y \leq 15$$

vertical line solid horiz dashed dashed

$$\textcircled{4} x + 6y \leq 15$$

$$6y \leq -x + 15$$

$$y \leq -\frac{1}{6}x + \frac{5}{2}$$


solid

- right of $\textcircled{1}$

- above $\textcircled{2}$

- above $\textcircled{3}$

- below $\textcircled{4}$




Example 4 – Write the system of inequalities for the following solution set.

$$\textcircled{1} y \leq 3$$

$$\textcircled{2} y \geq 2x - 3$$

$$\textcircled{3} y \geq -x$$



word
problem

Example 5 – The Canucks have 8 games left to play and need 10 points to make the playoffs. A win is worth 2 points and an overtime loss is worth 1 point. Write and graph a system of linear inequalities to see all the possible ways the Canucks can make the playoffs.

Let x = number of wins

Let y = number of overtime losses

inequalities:

① $x \geq 0$ RIGHT ①

② $y \geq 0$ ABOVE ②


③ $x + y \leq 8$

④ $2x + y \geq 10$

rearranged:

③ $y \leq -\frac{1}{1}x + 8$ BELOW ③

④ $y \geq -\frac{2}{1}x + 10$ ABOVE ④



Ways to make the playoffs:

examples:

$(6, 1) \Rightarrow 6$ wins, 1 OT loss = 13 pts.

$(5, 0) \Rightarrow 5$ wins = 10 pts.

$(3, 4) \Rightarrow 3$ wins, 4 OT losses = 10 pts.

etc.


7.3 – Solving Quadratic Inequalities

In Grade 9, you learned to solve linear inequalities such as $x - 4 > 0$. One way to solve them was to isolate x , so in this case add 4 to both sides and you get: $x > 4$.

Another way to solve these is to use a graph:

$x - 4 > 0$
 notice 0 is in place of y ,
 and y is 0 on the x axis.

Graph $y = \frac{1}{1}x - 4$
 slope \uparrow y -int



$x - 4 > 0$
 Where is the line ABOVE the x axis?

The line is above the x axis at all x values greater than 4, or, $x > 4$.

We wouldn't regularly solve a linear inequality using a graph, because it is much easier to just solve it algebraically, as we did at first.

However, to solve a quadratic inequality, the graphing method is often the preferred method, as it is quicker. However, we will learn both the graphing method and the algebraic method.

Example 1 – Solve $x^2 + 2x > 8$ by graphing, and then using algebra (test intervals). Graph the solution on a number line.

Graphing

Steps


1. Get everything to the left side so that zero is on the right.
2. Find the roots (x -intercepts).
3. Sketch a graph and use the visual to solve the inequality.
 → if the quadratic is > 0 , find the domain where the graph is above the x -axis
 → if the quadratic is < 0 , find the domain where the graph is below the x -axis

1) $x^2 + 2x - 8 > 0$

2) Find the x -ints by temporarily changing the inequality symbol to equals.
 $x^2 + 2x - 8 = 0$ x -ints are $-4, 2$
 $(x+4)(x-2) = 0$

3) The a value is positive so the parabola opens up.

$x^2 + 2x - 8 > 0$
 Where is the parabola ABOVE the x -axis?
 Left of -4 and right of 2



Solution:

$x < -4, x > 2$

Solution on Number Line:



Algebraic (Test Interval) Steps

1. Find the critical numbers (the zeros) of the inequality.
2. Make an x-axis diagram of the resulting test intervals.
3. Test a value from each interval using the original inequality.

<p>1) $x^2 + 2x - 8 > 0$ $x^2 + 2x - 8 = 0$ $(x+4)(x-2) = 0$ x-ints (critical values): $-4, 2$</p>	<p>2) $\textcircled{\text{I}} x < -4$ $\textcircled{\text{II}} -4 < x < 2$ $\textcircled{\text{III}} x > 2$</p>	
<p>3) $\textcircled{\text{I}}$ Test -5 $(-5)^2 + 2(-5) - 8 > 0$ $25 - 10 - 8 > 0$ $7 > 0$ \checkmark</p>	<p>$\textcircled{\text{II}}$ Test 0 $0^2 + 2(0) - 8 > 0$ $-8 > 0$ \times</p>	<p>$\textcircled{\text{III}}$ Test 3 $(3)^2 + 2(3) - 8 > 0$ $9 + 6 - 8 > 0$ $7 > 0$ \checkmark</p>
<p>Solution: $x < -4, x > 2$</p>		

Example 2 – Solve $x^2 - 10x + 16 \leq 0$ using both methods and graph the solution on a number line


- *if the quadratic is ≥ 0 , find the domain where the graph is **above or on** the x-axis
- *if the quadratic is ≤ 0 , find the domain where the graph is **below or on** the x-axis

Graphing:

$x^2 - 10x + 16 \leq 0$
 \uparrow
 $a=1$ so opens up


x -ints
 $x^2 - 10x + 16 = 0$
 $(x-8)(x-2) = 0$
 x -ints are $2, 8$

$x^2 - 10x + 16 \leq 0$
 \uparrow
 Where is the parabola **BELOW** or **ON** the x -axis?
 between 2 and 8 , including 2 and 8



<p>Solution: $2 \leq x \leq 8$</p>	<p>Solution on Number Line:</p>
--	---------------------------------

Test Intervals:



$\textcircled{\text{I}}$ Test 0
 $0^2 - 10(0) + 16 \leq 0$
 $16 \leq 0$
 \times

$\textcircled{\text{II}}$ Test 5
 $5^2 - 10(5) + 16 \leq 0$
 $25 - 50 + 16 \leq 0$
 $-9 \leq 0$
 \checkmark

$\textcircled{\text{III}}$ Test 9
 $9^2 - 10(9) + 16 \leq 0$
 $81 - 90 + 16 \leq 0$
 $7 \leq 0$
 \times

Solution:
 $2 \leq x \leq 8$

Example 3 – Graph the quadratic function $f(x) = x^2 - 6x + 9$. What is the solution to:

a) $x^2 - 6x + 9 \geq 0$ b) $x^2 - 6x + 9 > 0$ c) $x^2 - 6x + 9 \leq 0$ d) $x^2 - 6x + 9 < 0$



$f(x) = x^2 - 6x + 9$
 $y = (x-3)^2$
 x-int at 3
 $a=1$ so opens up

ⓑ $x^2 - 6x + 9 > 0$

Where is the parabola ABOVE the x axis?

Everywhere except at $x=3$


Solution: $x \neq 3$



ⓐ $x^2 - 6x + 9 \geq 0$
 Where is the parabola ABOVE or ON the x axis?

Everywhere!

$x \in \mathbb{R}$



ⓒ $x^2 - 6x + 9 \leq 0$

Where is the parabola BELOW or ON the x axis?

Never Below, on only at $x=3$

Solution: $x=3$



ⓓ

$x^2 - 6x + 9 < 0$
 Where is the parabola BELOW the x axis?

It is never below!

Solution: \emptyset

Example 4 – Solve $x^2 - 2x > 2$. Then graph the solution on a number line.

$x^2 - 2x - 2 > 0$

$x^2 - 2x - 2 = 0$
 cannot factor so must use quad formula.


$$x = \frac{2 \pm \sqrt{(-2)^2 - 4(1)(-2)}}{2(1)}$$

$$x = \frac{2 \pm \sqrt{12}}{2}$$

$$x = \frac{2 \pm 2\sqrt{3}}{2}$$


x-ints are $x = 1 \pm \sqrt{3}$
 approx. $x = 2.73, -0.73$
 $a=1$ so opens up

$x^2 - 2x - 2 > 0$
 Where is the parabola ABOVE the x axis?
 Left of $1 - \sqrt{3}$ and right of $1 + \sqrt{3}$



Solution:
 $x < 1 - \sqrt{3}$,
 $x > 1 + \sqrt{3}$

Solution on Number Line:



7.5 – Applications of Systems & Systems of Inequalities

Example 1 – A certain website offers online interactive puzzles, but the puzzle-makers present the following problem for entry to their site. “Determine two integers such that the sum of the smaller number and twice the larger number is 46. Also, when the square of the smaller number is decreased by three times the larger, the result is 93. By determining the smaller and larger numbers, use it as a password to gain access to the site.

Let x = smaller number
Let y = larger number

$$\begin{array}{r} \textcircled{1} \quad x + 2y = 46 \\ \textcircled{2} \quad x^2 - 3y = 93 \end{array}$$

can solve this system using substitution or elimination

Elimination:

$$\begin{array}{r} \textcircled{2} \quad (x^2 - 3y = 93) \times 2 \\ \textcircled{1} \quad (x + 2y = 46) \times 3 \end{array}$$

$$\begin{array}{r} \textcircled{2} \quad 2x^2 - 6y = 186 \\ + \\ \textcircled{1} \quad (3x + 6y = 138) \\ \hline 2x^2 + 3x = 324 \\ 2x^2 + 3x - 324 = 0 \\ 2x^2 - 24x + 27x - 324 = 0 \\ 2x(x - 12) + 27(x - 12) = 0 \\ (x - 12)(2x + 27) = 0 \\ x = 12, \frac{-27}{2} \leftarrow \text{not an integer} \end{array}$$


$$\begin{array}{l} x = 12 \\ \textcircled{1} \quad 12 + 2y = 46 \\ 2y = 34 \\ y = 17 \end{array}$$

The password is
12 17
(12 is smaller integer,
17 is larger)

Example 2 – An automobile storage area can fit at most 100 cars & trucks on its lot. A car covers 100 sq feet and a truck 200sq ft of space on a lot that is 12 000 sq ft. What are all the possibilities of cars & trucks that can be on the lot at any one time?


Let x = number of cars
Let y = number of trucks

$$\begin{array}{l} \textcircled{1} \quad x \geq 0 \quad \text{RIGHT } \textcircled{1} \\ \textcircled{2} \quad y \geq 0 \quad \text{ABOVE } \textcircled{2} \\ \textcircled{3} \quad x + y \leq 100 \\ \textcircled{4} \quad 100x + 200y \leq 12000 \\ \textcircled{3} \quad y \leq -x + 100 \quad \text{BELOW } \textcircled{3} \\ \textcircled{4} \quad \frac{200y}{200} \leq \frac{-100x}{200} + \frac{12000}{200} \\ y \leq -\frac{1}{2}x + 60 \quad \text{BELOW } \textcircled{4} \end{array}$$



Example 3 - The height in metres of a projectile shot from the top of a building is given by $h(t) = -16t^2 + 60t + 25$, where t represents the time in seconds the projectile is in the air.

- a) Find the time the projectile is in the air before hitting the ground, to the nearest thousandth.
 b) Find the time interval that the projectile is above 25m, to the nearest hundredth.



a) $h(t) = 0$ when projectile hits the ground.

$$-16t^2 + 60t + 25 = 0$$

$$16t^2 - 60t - 25 = 0$$

$$t = \frac{60 \pm \sqrt{60^2 - 4(16)(-25)}}{32}$$

$$t = 4.128, -0.378$$

↑ reject

*Enrichment: Example 4 - The price a stereo will be sold for is given by $S(x) = 200 - 0.1x$,

$0 \leq x \leq 2000$, where x is the number of stereos produced each day. It costs \$18 000 per day to operate the factory and \$15 for material to produce each stereo.

- a) Find the daily revenue. (b) Find the daily cost of producing stereos. (c) Find the interval that produces a profit.

a) Revenue = (Items Sold)(Price)

$$R(x) = x(200 - 0.1x)$$

$$R(x) = -0.1x^2 + 200x$$

b) $C(x) = 15x + 18000$

c) $P(x) = R(x) - C(x)$

$$P(x) = -0.1x^2 + 200x - (15x + 18000)$$

$$P(x) = -0.1x^2 + 185x - 18000$$

The profit equation must be greater than 0 to turn a profit.

The projectile is in the air for 4.128 seconds


b) $-16t^2 + 60t + 25 > 25$

$$-16t^2 + 60t > 0$$

$$-4t(4t - 15) = 0$$

$$t = 0, \frac{15}{4}$$

a is negative so opens down



$-16t^2 + 60t > 0$
 Where is the parabola ABOVE the x axis?
 Between 0 and $\frac{15}{4}$

$$0 < t < \frac{15}{4}$$

$$0 < t < 3.75s.$$

The projectile is above 25m for the first 3.75s of its flight.

c) $-0.1x^2 + 185x - 18000 > 0$


Find x -ints

$$a = -0.1, b = 185, c = -18000$$

$$x = \frac{-185 \pm \sqrt{(185)^2 - 4(-0.1)(-18000)}}{2(-0.1)}$$

$$x = 103.04, 1746.96$$

$a = -0.1$ so opens down



Where is the parabola ABOVE the x axis?

Between 103.04 and 1746.96

x must be whole number so between 104 and 1746

$$104 \leq x \leq 1746$$

will produce a profit.

